



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number : **0 634 170 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number : **94304561.7**

(51) Int. Cl.⁶ : **A61K 31/47**

(22) Date of filing : **23.06.94**

(30) Priority : **05.07.93 GB 9313866**

(43) Date of publication of application :
18.01.95 Bulletin 95/03

(84) Designated Contracting States :
**AT BE CH DE DK ES FR GR IE IT LI LU MC NL
PT SE**

(71) Applicant : **DIOMED DEVELOPMENTS
LIMITED**
Tatmore Place, Gosmore
Nr. Hitchin
Hertfordshire, SG4 7QR (GB)

(72) Inventor : **Whitefield, Martin**
19 Parkway
London NW11 (GB)

(74) Representative : **Collier, Jeremy Austin Grey**
J.A.KEMP & CO.
14, South Square
Gray's Inn
London WC1R 5LX (GB)

(54) **Compositions containing 8-hydroxy-quinoline for use in the treatment of hyperproliferative skin diseases.**

(57) 8-Hydroxyquinoline is useful in the topical treatment of inflammatory proliferative skin diseases, especially psoriasis. It should be applied from an essentially anhydrous vehicle.

EP 0 634 170 A1

This invention relates to compositions for the topical treatment of inflammatory proliferative skin diseases, especially psoriasis, and their manufacture and use.

Psoriasis is associated with the rapid turnover of skin cells (hyperproliferation) accompanied by a loss of differentiation so that silvery white scales form on the surface of the skin. Additionally, the capillaries become tortuous and dilated and an inflammatory reaction occurs, so that the skin reddens. The elevated silvery white scales on a contrasting red background produce the unsightly lesions characteristic of psoriasis.

A number of methods are used for the topical treatment of psoriasis, but none is entirely satisfactory. Historically, the earliest form of therapy involved the use of coal tar derivatives. These are sometimes effective but are extremely messy. Subsequently, the use of dithranol (1,8-dihydroxy-9-anthrone) was developed. This substance is very effective, but it causes irritation of the skin and staining both of the skin and adjacent materials, which are disadvantages especially in home treatment. Topical corticosteroids have also been used. Although these are clean and convenient to use, they are only suppressive in action and therefore only provide short term relief of symptoms. In addition, psoriasis treated with corticosteroids tends to relapse in a more resistant form, so that more aggressive therapy is later required.

The latest form of topical treatment of psoriasis is the use of synthetic Vitamin D derivatives. These derivatives are reasonably effective but have to be used in extremely low concentrations in order to avoid systemic calcitropic effects. Thus, it has been shown that increasing the concentration of the derivative in the composition above 0.005% w/w is liable to cause an increase in plasma calcium levels and, even at the 0.005% w/w level used in the marketed licensed product, there have been reports of increased plasma calcium levels. Such preparations moreover rarely clear the psoriasis, perhaps because of the very low strength used. Patients therefore have to continue treatment for extended periods which increases the risk of systemic side effects. Furthermore, a significant proportion of patients experience irritation and this restricts usage, particularly on delicate areas of the skin such as the face. Although it is not known for certain, the Vitamin D appears to alter calcium metabolism in the epidermal cells, which results in reduced cell proliferation and increased cell differentiation, leading to a correction of the abnormal pathophysiology of the epidermal cells and a return to normal skin growth and development.

We have now found that 8-hydroxyquinoline is effective in the treatment of psoriasis and other inflammatory proliferative skin diseases by topical application to the affected area of skin. More particularly, we have found that 8-hydroxyquinoline at a concentration of 0.25% in a simple ointment or gel has a rapid ameliorating effect on patients with psoriasis. It rapidly removes the scales on resistant patches of psoriasis, such as those which occur on the elbows and knees.

A possible explanation for the effectiveness of 8-hydroxyquinoline may be that it deprives the epidermal cells of the calcium they need for rapid proliferation by chelating the calcium and thus rendering it unavailable. However other calcium chelating agents are not effective against psoriasis. For example ethylene diamine tetraacetic acid and its salts were tested in various formulations and found to have no clinical effect whatsoever. Similarly, other chelating agents, including penicillamine, glycine, dimercaprol, citric acid, and 5-chloro-7-iodo-8-hydroxyquinoline (the topical antiseptic sold under the registered Trade Mark "Vioform") were all found to have no beneficial effect in the topical treatment of psoriasis.

8-Hydroxyquinoline is a white or slightly off-white solid which is relatively insoluble in water but dissolves in many organic solvents. When dissolved in anhydrous ethanol at a concentration of 1% w/w, it produces a clear almost colourless solution. On addition of water, e.g. 20% of the total volume, the 8-hydroxyquinoline remains in solution but the colour changes to yellow. This implies that 8-hydroxyquinoline exists in different forms, depending on whether or not water is present, *viz* a yellow form in the presence of water and a colourless form under essentially anhydrous conditions.

Trials have been carried out with both aqueous systems (e.g. aqueous gels) and essentially anhydrous systems (e.g. ointments or anhydrous gels) and it has been found that only the essentially anhydrous systems are effective in the treatment of psoriasis. Small quantities of water, e.g. 3 to 4% by weight, can be present in the essentially anhydrous compositions without them becoming inactive, but the desired therapeutic effect is lost if more than a small amount of water is present. This loss of activity is associated with the development of the yellow colour shown by 8-hydroxyquinoline in the presence of water and the presence or absence of this yellow colour thus provides a convenient way of determining whether a composition contains too much water to be active. It is thus a very simple matter to determine by experiment how much water a particular formulation can tolerate before it becomes inactive.

Water is not the only substance capable of rendering 8-hydroxyquinoline unavailable and, in consequence, without useful therapeutic effect. More particularly, acids and bases render 8-hydroxyquinoline unavailable by either converting it into a salt or limiting its release by hydrogen bond formation. Also substances such as polyhydric alcohols, polyethylene glycol, polyethylene glycol ethers and esters and non-ionic surfactants in general prevent release of 8-hydroxyquinoline into the skin from the phase containing it. The presence of such

substances in the compositions of the invention must therefore be avoided. It is believed that all such substances which render 8-hydroxyquinoline therapeutically unavailable have the property already noted of converting it into a yellow form, a colour change which is believed to be associated with disturbance of the intramolecular hydrogen bonding of the 8-hydroxyquinoline, believed to be a prerequisite for efficient dermatological availability. The compositions of the invention are thus white (colourless) rather than yellow (unless a yellow colouring agent is added).

The present invention accordingly provides a pharmaceutical composition for topical application comprising 8-hydroxyquinoline in free form dissolved in a pharmaceutically acceptable essentially anhydrous solvent. Such compositions preferably comprise from 0.01 to 5.0% by weight, more preferably from 0.1 to 1.0% by weight, most preferably 0.1 to 0.5% by weight, of 8-hydroxyquinoline. The essentially anhydrous solvent is preferably paraffin or another topically acceptable, oleophilic, and water-immiscible solvent. A mixture of white soft paraffin and isopropyl myristate has the required properties and is preferred.

The pharmaceutical composition may further comprise a lipid soluble antioxidant which is chemically inert towards 8-hydroxyquinoline to protect the 8-hydroxyquinoline from atmospheric oxidation.

The pharmaceutical compositions may also comprise a topical corticosteroid, for example, betamethasone valerate, hydrocortisone, or another corticosteroid known to be suitable for topical application, in an amount of 0.01 to 1.0% by weight. These compositions are useful for treating cases where there is excessive inflammation present.

These pharmaceutical compositions are preferably for use in the treatment of an inflammatory proliferative skin disease, especially psoriasis.

The present invention further provides 8-hydroxyquinoline in free form for use in the treatment of, and in the manufacture of a medicament for the treatment of, an inflammatory proliferative skin disease, especially psoriasis, and a method of treating an inflammatory skin disease, especially psoriasis, which comprises applying 8-hydroxyquinoline in free form to skin affected by such disease.

Although the present invention is primarily concerned with the treatment of psoriasis, it is within the scope of the invention to use 8-hydroxyquinoline in the treatment of other inflammatory skin conditions in which hyperproliferation or thickening occurs, e.g. chronic eczema/dermatitis.

While a simple ointment containing 8-hydroxyquinoline in white soft paraffin is (as noted above) effective in the present invention, it is desirable to reduce the greasiness of the formulation while retaining the 8-hydroxyquinoline in an essentially anhydrous oleophilic environment. To achieve this objective, the compositions of the invention may be provided in the form of an oil-in-water semi-solid emulsion, in which the 8-hydroxyquinoline is dissolved in the discontinuous oil phase. The aqueous phase of such compositions must have a substantially neutral pH to ensure that the partition of 8-hydroxyquinoline is strongly in favour of the oil phase. In these circumstances, even though the continuous phase is aqueous, the 8-hydroxyquinoline is present in an essentially anhydrous environment because of the very low solubility of water in the dispersed oily phase. Such a composition may contain over 50% of a dispersed oil phase in which the 8-hydroxyquinoline is dissolved. Although the lipid phase is preferably preponderantly white soft paraffin, chosen for its inertness and water immiscibility, a small amount of isopropyl myristate is preferably added to improve emolliency. In addition, since 8-hydroxyquinoline may be subject to atmospheric oxidation, a small concentration of an antioxidant, may be included. It is also preferable that the aqueous phase contains a preservative, preferably phenoxyethanol in an amount of 0 to 0.2% by weight. This is because the 8-hydroxyquinoline, which is itself an antiseptic, is held almost entirely in the lipid phase.

In an alternative embodiment of the invention, the 8-hydroxyquinoline is provided as a therapeutically active gel by dissolving in it in free form in an essentially anhydrous pharmaceutically acceptable solvent and gelling with a thickener. For example, ethanol may be used which forms a clear colourless gel on the addition of hydroxypropyl cellulose. A small proportion of a soluble inert lipid, such as isopropyl myristate may be added to reduce the drying effect of the alcohol. Again, a small concentration of an antioxidant may be added. The amount of thickener to be added is adjusted to produce a final formulation varying from a slightly viscous liquid, for application as a lotion, to a semi-solid gel.

The following examples illustrate the invention. Parts are by weight.

EXAMPLE 1

	Most Preferred Range	Range
5 8-hydroxyquinoline	0.1, 0.25, 0.5 or 1.0	0.01 - 5.0
White Soft Paraffin BP	40.0	10 - 50
Isopropyl Myristate BP	10.0	0 - 30
10 Cetostearyl Alcohol BP	7.0	5 - 10
Sodium Lauryl Sulphate BP	1.0	0.5 - 2.0
Phenoxyethanol BP	0.1	0 - 0.2
15 Purified Water BP to	100.0	

The formulation is made as follows:

1. Melt together the White Soft Paraffin (40 parts), isopropyl myristate (10 parts) and Cetostearyl Alcohol (7 parts) by heating to 70°C.
- 20 2. Dissolve 8-hydroxyquinoline (0.5 parts) in the heated oily phase so obtained.
3. Boil water (41.35 parts) thoroughly to remove any dissolved oxygen and allow to cool to 70°C.
4. Dissolve Sodium Lauryl Sulphate (1 part) and Phenoxyethanol (0.1 part) in the water at 70°C.
5. Add the aqueous phase so obtained to the oily phase and homogenise continuously, avoiding aeration, allowing the mixture to cool until semi-solid, to produce an almost white cream.
- 25 6. Pack into aluminium collapsible tubes.

EXAMPLE 2

	Most Preferred Range	Range
30 8-hydroxyquinoline	0.1, 0.25, 0.5 or 1.0	0.01 - 2.0
35 Isopropyl myristate BP	10.0	5 - 15
40 Hydroxypropyl Cellulose* BP	3.0	0.5 - 3.5
45 Ethanol BP to	100.0	
50		

* - Klucel HF, Aqualon

The formulation is made as follows:

- 55 1. Dissolve the 8-hydroxyquinoline in the alcohol.
2. Add the isopropyl myristate and mix well.
3. Warm to 50-60°C.

4. Add the hydroxypropyl cellulose slowly with continuous mixing, avoiding aeration.
5. Allow to cool with occasional mixing to form a clear colourless gel.
6. Pack into a bottle or aluminium tube depending on the viscosity.

The preparation should be applied twice daily to the affected areas of psoriasis. For thin areas of psoriasis a concentration of 0.1% to 0.5% 8-hydroxyquinoline may be sufficient, while for the thicker areas a concentration of 0.5% or 1.0% 8-hydroxyquinoline may be required. The preparation is non-irritant, it produces no staining, and has no effect on systemic calcium levels. It has been shown to be at least as effective as Vitamin D derivatives in some subjects, and even more effective in others. 8-hydroxyquinoline is particularly useful in the treatment of resistant areas of psoriasis which may have to be treated for extended periods.

If desired, betamethasone valerate may be incorporated in the above formulations in a concentration of 0.1% by weight.

Claims

1. A pharmaceutical composition for topical application comprising 8-hydroxyquinoline in free form dissolved in a pharmaceutically acceptable essentially anhydrous solvent.
2. A pharmaceutical composition according to claim 1 wherein 8-hydroxyquinoline is present in amount of from 0.01 to 5.0% by weight.
3. A pharmaceutical composition according to claim 2 wherein 8-hydroxyquinoline is present in an amount of from 0.1 to 1.0% by weight.
4. A pharmaceutical composition according to claim 1, 2, or 3 wherein the said solvent is oleophilic and water-immiscible.
5. A pharmaceutical composition according to claim 4 wherein the essentially anhydrous oleophilic, water-immiscible solvent comprises a mixture of white soft paraffin and isopropyl myristate.
6. A pharmaceutical composition according to any one of claims 1 to 5 which also comprises an antioxidant.
7. A pharmaceutical composition according to any one of the preceding claims which also comprises a topical corticosteroid.
8. A pharmaceutical composition according to claim 7 wherein the topical corticosteroid is betamethasone valerate in an amount of from 0.01 to 1.0% by weight.
9. A pharmaceutical composition according to any one of claims 1 to 8 wherein the essentially anhydrous solution of 8-hydroxyquinoline is dispersed in an aqueous phase which has a substantially neutral pH.
10. A pharmaceutical composition according to claim 9 wherein the said aqueous phase contains an anionic dispersing agent.
11. A pharmaceutical composition according to claim 9 or 10 wherein 0 to 0.2% by weight of phenoxyethanol is dissolved in the aqueous phase.
12. A pharmaceutical composition according to claim 1, 2 or 3 in the form of a gel in which the 8-hydroxyquinoline in free form is dissolved in an essentially anhydrous pharmaceutically acceptable solvent gelled with a thickener.
13. A pharmaceutical composition according to claim 12 wherein the said solvent is ethanol and the said thickener is hydroxypropylcellulose.
14. A pharmaceutical composition according to any one of claims 1 to 13 for use in the treatment of an inflammatory proliferative skin disease.
15. A pharmaceutical composition according to claim 14 wherein the inflammatory proliferative skin disease is psoriasis.

16. Use of 8-hydroxyquinoline in free form in the manufacture of a medicament for the treatment of an inflammatory proliferative skin disease.

17. Use according to claim 16 wherein the medicament is as defined in any of claims 1 to 13.

5

18. Use according to claims 16 or 17 wherein the inflammatory proliferative skin disease is psoriasis.

10

15

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 30 4561

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	GB-A-981 144 (LICENCIA TALALMANYOKAT ERTEKESITO VALLALAT) * column 1, line 1 - column 2, line 73 * ---	1-4, 14, 15	A61K31/47
X	EP-A-0 278 660 (STIEFEL LABORATORIES)	1, 2, 4, 5, 9, 11, 14, 15	
Y	* page 3, line 46 - line 47; example 4 * ---	3, 6-8, 10, 12, 13, 16-18	
Y	US-A-4 038 388 (DAVID FREDERICK CLEAVER) * column 1, line 1 - column 2, line 34; example 2 * ---	3, 6-8, 10, 12, 13, 16-18	
Y	EUR. J. CANCER vol. 26, no. 8, 1990 pages 905 - 907 JARDENA NORDENBERG ET AL. 'Anti-proliferative effects and phenotypic alterations induced by 8-hydroxyquinoline in melanoma cell lines' * abstract * ---	3, 6-8, 10, 12, 13, 16-18	TECHNICAL FIELDS SEARCHED (Int.Cl.6) A61K
Y	DERMATOLOGY NURSING vol. 3, no. 1, 1991 pages 25 - 28 SHARON M. MINNICH ET AL. 'effect of oxyquinoline ointment on diaper dermatitis' * abstract * -----	3, 6-8, 10, 12, 13, 16-18	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 12 September 1994	Examiner Tzschoepe, D
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 150 (03.92) (P4/C61)